



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Contents lists available at ScienceDirect

American Journal of Emergency Medicine

journal homepage: [www.elsevier.com/locate/ajem](http://www.elsevier.com/locate/ajem)

# ST-elevation in an adolescent with COVID-19: Myopericarditis or myocardial infarction?

Cathie-Kim Le, MD<sup>a,\*</sup>, Minh B. Nguyen, MD<sup>b</sup>, Andrea T. Vo, MD<sup>a</sup>

<sup>a</sup> Department of Emergency Medicine, Hospital for Sick Children, Toronto, Canada

<sup>b</sup> Department of Cardiology, Hospital for Sick Children, Toronto, Canada

## ARTICLE INFO

### Article history:

Received 9 June 2021

Received in revised form 19 August 2021

Accepted 24 August 2021

Available online xxxx

### Keywords:

COVID-19

Multisystem inflammatory syndrome in children

MIS-C

Pericarditis

Myocarditis

Myopericarditis

Myocardial infarction

## ABSTRACT

Myocardial infarctions (MI) have been reported in adults with COVID-19. Although MIs are rare in children with COVID-19, cardiac involvement is still possible. In this case report, we present an adolescent with recent COVID-19 infection who presented with an ECG initially suggestive of myocardial infarction (MI). We describe how to differentiate between myocardial infarctions and myopericarditis. A 15-year-old boy, with a history of COVID-19 infection a month prior, presented to the emergency department with fever, abdominal pain, diarrhea, and chest pain. On ECG, he was found to have focal ST-segment elevations in V3 through V6. Given the immediate concern for MI, an emergent echocardiogram was done and showed normal left ventricular systolic function with no regional dyskinesia and normal coronary artery diameters. A repeat ECG showed diffuse ST elevations in the inferior leads and T-wave inversions on V5 and V6, confirming the diagnosis of myopericarditis. In conclusion, multisystem-inflammatory syndrome in children associated with COVID-19 (MIS-C) is a new entity describing a post-infectious inflammatory response in children with prior COVID-19 exposure. Cardiac involvement can include myopericarditis. Initial ECGs may show ST-changes suggestive of MI. However, serial ECGs and echocardiograms can differentiate between MI and myocarditis/myopericarditis. Even with COVID-19, MIs are extremely rare in children, and it is important to be aware of MIS-C and its cardiac complications.

Crown Copyright © 2021 Published by Elsevier Inc. All rights reserved.

## 1. Background

Myocardial infarctions (MIs) have been reported in adults with COVID-19 but remain extremely rare in children. We present a case of a patient who presented with ECG findings initially suggestive of MI but was ultimately diagnosed with myopericarditis associated with Multisystem Inflammatory Syndrome in Children (MIS-C), a new post-infectious inflammatory complication of COVID-19 exposure. We describe how to differentiate between MI and myopericarditis in patients with MIS-C.

## 2. Case presentation

A 15-year-old male, who had COVID-19 infection a month prior, presented in the emergency department with 4 days of fever, abdominal

pain, diarrhea, and chest pain. He had normal vital signs. Investigations revealed a CRP of 287.1 mg/L (normal 0.1–1.7 mg/L), an ESR of 64 mm/h (normal 2–28 mm/h), thrombocytopenia to  $106 \times 10^9/L$ , lymphopenia to  $0.46 \times 10^9/L$  and an elevated ferritin of 756.8 µg/L (normal 11.1–171.9 µg/L).

His troponin I was 1.74 ng/mL (normal <0.04 ng/mL) and his N-terminal pro-brain natriuretic peptide (NT-proBNP) was 1458 pg/mL (normal <125.0 pg/mL). Initial ECG (Fig. 1) revealed 3–5 mm ST segment elevations in V3 through V6. Because of concerns for MI, the patient was given aspirin, sublingual nitroglycerin, and placed on supplemental oxygen after cardiology consultation. An emergent echocardiogram showed good left ventricular systolic function, no regional dyskinesia and normal coronary artery diameter. Since the echocardiogram was normal, an acute coronary event was deemed unlikely, and the patient was admitted with the presumptive diagnosis of myopericarditis secondary to MIS-C. Repeat ECG (Fig. 2) the next day revealed a reduction of ST elevation in the anterolateral precordial leads, but now with >1 mm ST elevation in the inferior leads. In addition, the T waves in V5 and V6 had become inverted.

He was treated with IV methylprednisolone, IV immunoglobulin, and daily aspirin. Repeat echocardiogram on day 3 showed a small left anterior descending coronary aneurysm, normally sized right coronary artery, and good ventricular systolic function. The patient was discharged on day 3 of hospitalization with no residual cardiac or

**Abbreviations:** CRP, C-reactive protein; ESR, erythrocyte sedimentation rate; MI, myocardial infarction; MIS-C, Multisystem inflammatory syndrome in children associated with COVID-19; NT-pro-BNP, N-terminal pro-brain natriuretic peptide; STEMI, ST-elevation myocardial infarction.


\* Corresponding author at: Department of Emergency Medicine, 525 University Avenue, Toronto M5G 2L3, Canada.

E-mail address: [cathie-kim.le@sickkids.ca](mailto:cathie-kim.le@sickkids.ca) (C.-K. Le).

<https://doi.org/10.1016/j.ajem.2021.08.068>

0735-6757/Crown Copyright © 2021 Published by Elsevier Inc. All rights reserved.

Please cite this article as: C.-K. Le, M.B. Nguyen and A.T. Vo, ST-elevation in an adolescent with COVID-19: Myopericarditis or myocardial infarction?, American Journal of Emergency Medicine, <https://doi.org/10.1016/j.ajem.2021.08.068>



**Fig. 1.** Initial ECG showing ST segment elevations in V3 through V6.

gastrointestinal symptoms and a prescription for prednisolone taper and aspirin, with cardiology follow-up.

A week later, the patient's symptoms returned to baseline. Close follow-up with cardiology revealed improving coronary artery diameters that ultimately normalized. ECG showed normal sinus rhythm with non-specific ST-segment and T-wave changes. The patient was discharged from cardiology five months later with no cardiac sequelae.

### 3. Discussion


Although ST elevation myocardial infarction (STEMI) has been described in adults with COVID-19, only one case has been reported in children [1]. There is, however, a growing number of children who present with myopericarditis in the context of MIS-C. [2,3] The hallmark feature of MIS-C is high persistent fevers for  $\geq 3$  days with features of marked systemic inflammation and a temporal association with COVID-19 present in the community 3–6 weeks prior to presentation [4]. The pathophysiology of MIS-C is thought to be a post-infectious inflammatory process related with COVID-19 infection [4].

Pericarditis is inflammation of the pericardium. Although the pericardium is electrically silent, ECG anomalies can be found secondary to pericardial effusion or inflammation of the adjacent superficial myocardium [5]. ECG findings typically show diffuse concave ST elevations [6]. Myocarditis is inflammation of the myocardium in the absence of ischemia and may demonstrate nonspecific ST segment abnormalities, T-waves abnormalities, conduction abnormalities or decreased QRS

voltage [6]. ECG changes in MIS-C are most consistent with changes seen in myopericarditis: low amplitude ECGs, transient T-wave inversion in lateral precordial leads, either non-specific or diffuse evolving ST changes, and rarely AV block.[1,7] Compared to MI, ST-segment changes in myopericarditis tend to be less pronounced, rarely exceeding 4–5 mm. In addition, those changes tend to be seen on multiple leads: I, II, V5 and V6 with reciprocal changes in aVR and V1. Furthermore, in myopericarditis, T-wave changes are usually present in all leads except aVR and V1, and are less deep or incompletely inverted. Finally, ECG changes in myopericarditis tend to happen more slowly and recover faster compared to MI [5].

Our patient had an initially concerning ECG for MI, but the evolution of diffuse ST elevations and T wave inversions on subsequent ECG indicated myopericarditis. On echocardiogram, STEMI generally will reveal focal wall motion abnormality, whereas myocarditis will likely demonstrate either normal wall motion or global systolic dysfunction [8]. Coronary artery aneurysms or dilatations may also be present, in addition to elevated levels of BNP and troponin [8]. Our patient's echocardiogram findings therefore supports the final diagnosis of myopericarditis rather than a true ischemic event.

Myocarditis from MIS-C can be associated with significant morbidity including multi-organ failure, heart failure and arrhythmias [2]. Treatment includes immunomodulatory drugs, including IV immunoglobulin and systemic glucocorticoids [9,10]. In most cases, the patient's ventricular function normalizes but the long-term effect remains unclear [11].



**Fig. 2.** Repeat ECG showing diffuse ST elevation in anterolateral precordial leads and inferior leads.

In conclusion, children presenting with myopericarditis in the context of MIS-C may show ST-changes suggestive of STEMI. Serial ECGs and echocardiograms can differentiate between MI and myocarditis/myopericarditis. It is important to recognize MIS-C and its cardiac complications among children to initiate the appropriate treatment and management.

### Declaration of Competing Interest

none.

### References

- [1] Regan W, O'Byrne L, Stewart K, et al. Electrocardiographic changes in children with multisystem inflammation associated with COVID-19. *J Pediatr*. 2021;234:27–32. <https://doi.org/10.1016/j.jpeds.2020.12.033>.
- [2] Belhadj Z, Méot M, Bajolle F, et al. Acute heart failure in multisystem inflammatory syndrome in children in the context of global SARS-CoV-2 pandemic. *Circulation*. 2020;142(5):429–36. <https://doi.org/10.1161/CIRCULATIONAHA.120.048360>.
- [3] Grimaud M, Starck J, Levy M, et al. Acute myocarditis and multisystem inflammatory emerging disease following SARS-CoV-2 infection in critically ill children. *Ann Intensive Care*. 2020;10(1):69. <https://doi.org/10.1186/s13613-020-00690-8>.
- [4] Berard RA, Haddad EM, Morin MP, Chan K, Dahdah NS, McCrindle BW, et al. Paediatric inflammatory multisystem syndrome temporally associated with COVID-19. *Canadian Paediat Soc*. July 2020 <https://www.cps.ca/documents/position/pims>.
- [5] Surawicz B, Lasserter KC, FCD, DoM, UoK, CoMLKUSA. Electrocardiogram in pericarditis. *Am J Cardiol*. 1970;26(5):471–4. [https://doi.org/10.1016/0002-9149\(70\)90704-6](https://doi.org/10.1016/0002-9149(70)90704-6).
- [6] Allen HD, Shaddy RE, Penny DJ, et al. Moss and Adams' heart disease in infants, children, and adolescents : Including the fetus and young adult. Wolters Kluwer: Ninth edition. ed. Philadelphia; 2016.
- [7] Choi NH, Fremed M, Starc T, et al. MIS-C and cardiac conduction abnormalities. *Pediatrics*. 2020;146(6):e2020009738. <https://doi.org/10.1542/peds.2020-009738>.
- [8] Feldstein LR, Rose EB, Horwitz SM, et al. Multisystem Inflammatory Syndrome in U.S. Children and Adolescents. *N Engl J Med*. 2020;383(4):334–46. <https://doi.org/10.1056/NEJMoa2021680> [published Online First: 2020/07/01].
- [9] Long B, Brady WJ, Koyfman A, et al. Cardiovascular complications in COVID-19. *Am J Emerg Med*. 2020;38(7):1504–7. <https://doi.org/10.1016/j.ajem.2020.04.048> [published Online First: 2020/04/23].
- [10] Alsaied T, Aboulhosn JA, Cotts TB, et al. Coronavirus Disease 2019 (COVID-19) Pandemic Implications in Pediatric and Adult Congenital Heart Disease. *J Am Heart Assoc*. 2020;9(12). <https://doi.org/10.1161/jaha.120.017224> e017224. [published Online First: 2020/05/23].
- [11] Kaushik S, Aydin SI, Derespina KR, et al. Multisystem Inflammatory Syndrome in Children Associated with Severe Acute Respiratory Syndrome Coronavirus 2 Infection (MIS-C): A Multi-institutional Study from New York City. *J Pediatr*. 2020;224:24–9. <https://doi.org/10.1016/j.jpeds.2020.06.045> [published Online First: 06/14].